
**Effect of contact interactions on higgs production
cross-section at an e^+e^- collider**

**Uma Mahanta
Mehta Research Institute
Chhatnag Road, Jhusi
Allahabad-221506, India**

Abstract

New interactions appearing at a scale Λ larger than the weak interaction scale v can affect physical processes at energies below Λ through non-renormalizable $SU(3) \times SU(2) \times U(1)_y$ invariant operators added to the standard model Lagrangian. In this article we investigate the effect of flavor conserving contact interactions on the total cross-section for the process $e^+e^- \rightarrow HZ$ at $\sqrt{s} = 500$ Gev. We find that for $\Lambda \approx 2.5$ Tev, which is consistent with LEP and SLD asymmetry measurements on Z peak as well as theoretical estimates, these operators can increase the total cross-section by a factor of 3 relative to the SM for intermediate mass higgs boson.

The SM has been extremely successful in explaining all the experimental data so far. However in spite of its extraordinary phenomenological success many theorists regard the SM as an effective low energy theory valid below some cut off scale Λ of the order of a few Tev. One of the fundamental reasons behind this view is that the Higgs mass in the SM receives radiative corrections that diverges quadratically with the cut off Λ [1]. Hence to stabilize the higgs mass around the weak scale, which is the natural upper bound for m_H , the cut off scale Λ should be of the order of a few Tev. New interactions can appear at or above the scale Λ involving new heavy particles. Their effects on physical processes at energies below Λ can be described by an effective Lagrangian containing $SU(3)_c \times SU(2) \times U(1)_y$ invariant non-renormalizable operators involving only the light SM fields [2]. Since we shall be considering the effects of Tev scale new physics on higgs production cross-section the light SM fields should include ϕ . For the same reason the gauge symmetry will be assumed to be linearly realized on the SM fields. In addition to gauge symmetries one might also impose other constraints like baryon number and lepton number conservation. The non-renormalizable operators can be expressed as a systematic power series expansion in $\frac{1}{\Lambda}$. The lowest dimensional operator will clearly have the most dominant effect at energies below Λ . Further the effect of these non-renormalizable operators increases as the characteristic energy scale of the process under study approaches Λ .

In this article we shall consider the effect of flavor conserving d=6 operators involving leptons, scalar and gauge fields on intermediate mass ($m_H \approx 100\text{-}250$ Gev) higgs production via the process $e^+e^- \rightarrow HZ$ at $\sqrt{s} = 500$ Gev. We find that for $\Lambda \approx 2.5$ Tev the flavor conserving operators satisfy the LEP and SLD constraints on A_{fb}^e and A_{LR} . However the same value of Λ can increase the cross-section (if the new physics contribution interferes constructively with that of SM) for the process by a factor of 3 relative to that of the SM for a higgs mass of 150-200 Gev. On the other hand for destructive interference the cross-section decreases by a factor of .7 relative to SM for the same range of values of Λ

and m_H . The production and detection of intermediate higgs will be one of the important tasks of future e^+e^- collider. The detection of such a higgs boson will be extremely difficult at hadron collider since the higgs decays mainly into $b\bar{b}$ which can remain hidden in the background arising from $t\bar{t}$ pair production [3]. It is therefore extremely important to consider the effects of Tev scale new physics on higgs production cross-section at future e^+e^- colliders.

The effective Lagrangian is given by $L = L_{SM} + L_\Lambda$ where $L_\Lambda = \sum_i C_i O_i$. The coefficient C_i is of the order of Λ^{-2} where Λ is the characteristic scale for the operator O_i . The operators O_i of d=6 that can contribute to the process $e^+e^- \rightarrow HZ$ are [2]

$$\begin{aligned} O_1 &= (\bar{l}\gamma^\mu l) \frac{i}{2} [\Phi^+ (D_\mu \Phi) - (D_\mu \Phi)^+ \Phi] \\ &= \frac{v^2 g}{4c_w} \bar{e}_L \gamma^\mu e_L Z_\mu + \frac{vg}{2c_w} \bar{e}_L \gamma^\mu e_L Z_\mu H + \end{aligned} \quad (1)$$

$$\begin{aligned} O_2 &= (\bar{e}\gamma^\mu e) \frac{i}{2} [\Phi^+ (D_\mu \Phi) - (D_\mu \Phi)^+ \Phi] \\ &= \frac{v^2 g}{4c_w} \bar{e}_R \gamma^\mu e_R Z_\mu + \frac{vg}{2c_w} \bar{e}_R \gamma^\mu e_R Z_\mu H + \end{aligned} \quad (2)$$

$$\begin{aligned} O_3 &= (\bar{l}\gamma^\mu \tau_a l) \frac{i}{2} [\Phi^+ \tau_a (D_\mu \Phi) - (D_\mu \Phi)^+ \tau_a \Phi] \\ &= \frac{v^2 g}{4c_w} \bar{e}_L \gamma^\mu e_L Z_\mu + \frac{vg}{2c_w} \bar{e}_L \gamma^\mu e_L Z_\mu H + \end{aligned} \quad (3)$$

$$\begin{aligned} O_4 &= (\bar{l} D^\mu e) D_\mu \Phi + h.c. = \frac{igv}{2\sqrt{2}c_w} [(\partial_\mu \bar{e}_R) e_L - \bar{e}_L (\partial_\mu e_R)] Z^\mu \\ &+ \frac{ig}{2\sqrt{2}c_w} [(\partial_\mu \bar{e}_R) e_L - \bar{e}_L (\partial_\mu e_R)] H Z^\mu \\ &+ \frac{igs_w^2}{\sqrt{2}c_w} (\bar{e}_L e_R - e_L \bar{e}_R) Z^\mu \partial_\mu H + \frac{ie}{\sqrt{2}} (\bar{e}_R e_L - \bar{e}_L e_R) A^\mu \partial_\mu H + \end{aligned} \quad (4)$$

$$\begin{aligned}
O_5 &= (D^\mu \bar{l}) e D_\mu \Phi + h.c. = \frac{igv}{2\sqrt{2}c_w} [\bar{e}_R (\partial_\mu e_L) - (\partial_\mu \bar{e}_L) e_R] Z^\mu \\
&+ \frac{ig}{2\sqrt{2}c_w} [\bar{e}_R (\partial_\mu e_L) - (\partial_\mu \bar{e}_L) e_R] H Z^\mu \\
&+ \frac{ig(c_w^2 - s_w^2)}{2\sqrt{2}c_w} (\bar{e}_L e_R - \bar{e}_R e_L) Z^\mu \partial_\mu H + \frac{ie}{\sqrt{2}} (\bar{e}_L e_R - \bar{e}_R e_L) A^\mu \partial_\mu H + \dots
\end{aligned} \tag{5}$$

$$\begin{aligned}
O_6 &= (\bar{l} \sigma^{\mu\nu} \tau_3 e) \phi W_{\mu\nu}^3 + h.c. \\
&= -\frac{1}{\sqrt{2}} (\bar{e}_L \sigma^{\mu\nu} e_R) H (s_w F_{\mu\nu} + c_w Z_{\mu\nu}) + h.c. + \dots
\end{aligned} \tag{6}$$

$$\begin{aligned}
O_7 &= (\bar{l} \sigma^{\mu\nu} e) \phi B_{\mu\nu} + h.c. \\
&= \frac{1}{\sqrt{2}} (\bar{e}_L \sigma^{\mu\nu} e_R) H (c_w F_{\mu\nu} - s_w Z_{\mu\nu}) + h.c. + \dots
\end{aligned} \tag{7}$$

In the above we have expressed ϕ in unitary gauge and have written only the resulting d=4, 5 and 6 operators involving charged lepton, scalar and neutral gauge fields that can contribute to precision measurements on Z peak and to $e^+e^- \rightarrow HZ$. On transforming the lepton fields from the gauge eigenstate basis to the mass eigenstate basis the operators O_6 & O_7 give rise to d=5 FCNC operators involving both Z_μ and A_μ . In particular the coefficients C_6 and C_7 must satisfy the strong experimental bound [4] on the branching fraction for the process $\mu \rightarrow e\gamma$ which implies that

$$\frac{\Gamma_{\mu \rightarrow e\gamma}}{\Gamma_{\mu \rightarrow e\bar{\nu}_e \nu_\mu}} \approx \frac{6\pi^2 v^6}{\Lambda^4 m_\mu^2} (c_w - s_w)^2 \sin^2 \theta_{12} \leq 5 \times 10^{-11}. \tag{8}$$

If we assume that $\sin \theta_{12} \approx .2$, the scale $\Lambda \approx C_6^{-\frac{1}{2}} \approx C_7^{-\frac{1}{2}}$ associated with O_6 and O_7 must be greater than 3500 Tev. Such a huge scale for O_6 and O_7 can be avoided by assuming a symmetry that forbids FCNC vertices upon transformation from the gauge

basis to the mass basis. However the flavor diagonal terms of O_6 and O_7 must still satisfy the constraint [5] arising from the experimental value of $\frac{g_e-2}{2}$ which implies that

$$[(\frac{\delta g_e}{2})_{expt} - (\frac{\delta g_e}{2})_{sm}] = (\frac{\delta g_e}{2})_{new} \approx \frac{2\sqrt{2}m_e v(c_w - s_w)}{\Lambda^2 e} \leq .27 \times 10^{-9}. \quad (9)$$

Hence the scale Λ associated with the flavor diagonal terms of O_6 and O_7 must be greater than or of the order of 40 Tev which is considerably greater than the bound (2.5 Tev) on the scale associated with O_1 , O_2 and O_3 that follows from Z pole precision measurements. We shall therefore ignore effect of O_6 & O_7 on the process $e^+e^- \rightarrow HZ$. The remaining operators O_i , $i=1-5$ do not contain any $ee\gamma$ vertex but they do contain eeZ vertex. The scale associated with these flavor diagonal operators are best constrained by precision measurements on Z pole. However O_1 , O_2 and O_3 affect Z pole physics through d=4 operators, but O_4 & O_5 contribute to the same through d=5 operators. Hence the constraint on C_4 and C_5 that follows from Z pole precision measurements is expected to be weaker than that on C_1 , C_2 and C_3 . If we assume that $C_4 = C_5 = \frac{1}{\Lambda^2}$ we get

$$C_4 O_4 + C_5 O_5 \approx \frac{iC_4 g}{\sqrt{2}c_w} (\bar{e}\gamma_5 e) Z^\mu \partial_\mu H. \quad (10)$$

where we have integrated by parts and have used the relation $\partial_\mu Z^\mu = 0$ for on shell Z boson. Note that O_4 and O_5 contributes to Z pole precision measurements through d=5 terms in contrast to $O_1 - O_3$ which contributes to the same via d=4 terms. Hence the constraint on C_4 and C_5 that follows from Z pole precision measurements is expected to be slightly weaker than that on C_1 , C_2 and C_3 . In this work we shall determine the scale associated with O_1 , O_2 and O_3 from the LEP and SLD data and equate it to the scale associated with O_4 and O_5 . The effective Lagrangian that contributes to the process $e^+e^- \rightarrow HZ$ becomes

$$L_{eff} = \frac{g}{2c_w} [(1 - 2s_w^2) + \frac{v^2}{2}(C_1 + C_3)] \bar{e}_L \gamma^\mu e_L Z_\mu + \frac{g}{2c_w} [\frac{v^2}{2}C_2 - 2s_w^2] \bar{e}_R \gamma^\mu e_R Z_\mu$$

$$\begin{aligned}
& + \frac{vg^2}{4c_w^2} H Z_\mu Z^\mu + (C_1 + C_3) \frac{vg}{2c_w} \bar{e}_L \gamma^\mu e_L Z_\mu \\
& + C_2 \frac{vg}{2c_w} \bar{e}_R \gamma^\mu e_R Z_\mu + \frac{iC_4 g}{\sqrt{2}c_w} \bar{e} \gamma_5 e Z^\mu \partial_\mu H.
\end{aligned} \tag{11}$$

To a first approximation we can neglect the small corrections (of the order of 1%) to the $Ze\bar{e}$ vertex due to new physics. The total cross-section for $e^+e^- \rightarrow HZ$ is given by

$$\begin{aligned}
\sigma_T &= \sigma_{LR+RL} + \sigma_{LL+RR} \\
&= \frac{1}{384\pi s} \frac{f(s, M_z, M_H)}{s} \left[\left\{ \frac{2\pi\alpha}{s_w^2 c_w^2} \frac{(1 - 2s_w^2)}{(s - M_z^2)} + (C_1 + C_3) \right\}^2 \right. \\
&\quad \left. + \left\{ C_2 - \frac{2\pi\alpha}{s_w^2 c_w^2} \right\}^2 [12sM_z^2 + f^2(s, M_z, M_H)] \right] \\
&\quad + \frac{1}{64\pi v^2} \frac{f(s, M_z, M_H)}{s} C_4^2 f^2(s, M_z, M_H).
\end{aligned} \tag{12}$$

where $f^2(s, M_z, M_H) = (s + M_z^2 - M_H^2)^2 - 4sM_z^2$. The first term arises from $e_L^- e_R^+ + e_R^- e_L^+ \rightarrow HZ$ and the second term from $e_L^- e_L^+ + e_R^- e_R^+ \rightarrow HZ$. We shall consider two distinct cases of new physics effects on σ_T . In the first scenario we shall assume that new physics effects on σ_{LR+RL} interferes constructively with those of SM. A typical example of this scenario is $C_1 = C_3 = C_4 = -\frac{C_2}{2} = \frac{1}{\Lambda^2}$. In the second scenario the new physics effects on σ_{LR+RL} will be assumed to interfere destructively with SM effects, an example of which is $C_1 = C_3 = -C_4 = -\frac{C_2}{2} = -\frac{1}{\Lambda^2}$. We shall assume a common value of $\Lambda=2.5$ Tev for both scenarios.

The operators O_1 , O_2 and O_3 give rise to small corrections to the vector and axial vector couplings of Z boson to $e\bar{e}$. It is therefore important to consider the constraints on C_1 , C_2 and C_3 that follow from LEP and SLD asymmetry measurements on Z peak and whether these constraints are consistent with the value of Λ assumed above. We have $g_v^e = (g_v^e)_{sm} - \frac{v^2}{8}(C_1 + C_2 + C_3)$ and $g_a^e = (g_a^e)_{sm} + \frac{v^2}{8}(C_1 + C_3 - C_2)$. Hence in both scenarios $\delta g_v^e \approx 0$ and $\delta g_a^e \approx -\frac{v^2}{4}C_2$ i.e. new physics effects renormalizes the weak axial charge of the electron but does not affect its weak vector charge. On Z peak [6] we have $A_{LR} \approx (A_{LR})_{sm} [1 - \frac{\delta g_a^e}{(g_a^e)_{sm}}]$ and $A_{fb}^e \approx (A_{fb}^e)_{sm} [1 - 2\frac{\delta g_a^e}{(g_a^e)_{sm}}]$ provided $|\frac{\delta g_v^e}{(g_v^e)_{sm}}| \ll$

$|\frac{\delta g_a^e}{(g_a^e)_{sm}}|$. It then follows that $\frac{(\delta A_{LR})_{new}}{(A_{LR})_{sm}} \approx \pm \frac{v^2}{2\Lambda^2(g_a^e)_{sm}}$ and $\frac{(\delta A_{fb}^e)_{new}}{(A_{fb}^e)_{sm}} \approx \pm \frac{v^2}{\Lambda^2(g_a^e)_{sm}}$ where the upper (lower) sign corresponds to destructive (constructive) interference scenario. The average values of A_{LR} and A_{fb}^e reported by SLD and LEP [7] are $A_{LR} = .156 \pm .008$ and $A_{fb}^e = .0160 \pm .0024$. The predicted values for these asymmetries in the context of the SM are $(A_{LR})_{sm} = .142 \pm .003 \pm .003$ and $(A_{fb}^e)_{sm} = .0151$. The experimental bounds on δA_{LR} and δA_{fb}^e are $\frac{\delta A_{LR}}{(A_{LR})_{sm}} \approx .099$ and $\frac{\delta A_{fb}^e}{(A_{fb}^e)_{sm}} \approx .06$. For $\Lambda \approx 2.5$ Tev the new physics contributions are given by $\frac{\delta A_{LR}}{(A_{LR})_{sm}} \approx \pm .02$ and $\frac{\delta A_{fb}^e}{(A_{fb}^e)_{sm}} \approx \pm .04$ both of which are clearly compatible with the experimental bounds. On the contrary the contribution of O_4 to A_{LR} at the Z pole is given by $\frac{\delta A_{LR}}{(A_{LR})_{sm}} \approx -\frac{1}{2} \frac{(\frac{g}{4c_w})^2 \frac{v^2 s}{\Lambda^4}}{(g_L^e)^2 + (g_R^e)^2}$. The current SLD precision for measuring A_{LR} is about 5%. Hence even if Λ is as low as 400 Gev the contribution of O_4 to A_{LR} is far too small (about .78%) to be detected at SLD. In the following we shall equate the scale of O_4 and O_5 to that of O_1 , O_2 and O_3 . This will give us a lower bound on the cross-section for the process $e^+e^- \rightarrow HZ$. Further if the positive sign of the phenomenological bounds on δA_{LR} and δA_{fb}^e is taken seriously, experiments would seem to prefer the destructive interference scenario considered in this article, implying that the observed higgs production cross-section will be lower than that of the SM.

From eqn. 11 we find that for $\sqrt{s} = 500$ Gev and $m_H = 150$ Gev, $\sigma_T \approx 160$ fb (43 fb) for the constructive (destructive) interference case. This is to be compared with the SM prediction of $\sigma_{sm} \approx 52$ fb. When m_H is increased to 200 Gev, σ_T drops to 128 fb in the constructive case and to 33 fb in the destructive case. The corresponding value of the cross-section in the context of SM is 42 fb. In the case of constructive interference the dominant contribution to σ_T comes from σ_{LR+RL} whereas for destructive interference σ_{LL+RR} forms the dominant part. σ_T is therefore quite insensitive to the value of C_4 in the former case but depends quite strongly on it in the latter. For an integrated luminosity of 30 fb^{-1} [8] the effect of new physics would be to increase the number of HZ events by 3060 for constructive interference or decrease it by 390 for destructive interference. Note that at low energies ($\sqrt{s} \approx 500$ Gev) the dominant production mechanism for intermediate

mass higgs boson at an e^+e^- collider is $e^+e^- \rightarrow HZ$ [8]. For higher energies ($\sqrt{s} \approx 1$ Tev) the dominant production mechanism becomes $e^+e^- \rightarrow \nu_e \bar{\nu}_e H$. However new physics affects the latter process only through small corrections (of the order of 1%) to the usual SM vertices. There is no d=5 or 6 operator for the vertex $\bar{e}e\bar{\nu}_e\nu_e H$ similar to the $\bar{e}eZH$ vertex. Hence the overall effect of Tev scale new physics on the process $e^+e^- \rightarrow \nu_e \bar{\nu}_e H$ is expected to be much smaller than that on $e^+e^- \rightarrow HZ$.

Some comments are in order about the reliability of the value of Λ , the scale of new physics, used in our analysis. The scale Λ for new interactions should be related [9] to the amplitude v (v plays the role of f_π in electro-weak theory) for producing scalar particles out of the vacuum through the relation $\frac{\Lambda}{v} = g_s$ where g_s is the induced coupling for the low energy theory. According to theoretical estimates g_s is expected to lie between 1 and 4π . In low energy QCD for example $g_\rho = \frac{M_\rho}{f_\pi} \approx 6$. For $\Lambda \approx 2.5$ Tev we find that $g_s \approx 10$ which is in agreement with our theoretical expectations about new interactions underlying the EW theory.

References

1. The quadratic instability of the higgs sector was first pointed out by K. Wilson, unpublished. E. Gildener, Phys. Rev. D 14, 1667 (1976); L. Susskind, Phys. Rev. D 20, 2619 (1979).
2. W. Buchmuller and D. Wyler, Nucl. Phys. B 268, 621 (1986).
3. J. Gunion, S. Dawson, H. Haber and G. Kane, "The Higgs Hunters Guide", Addison Wesley (Menlo-Park, 1991).
4. Review of Particle Properties, Phys. Rev. D 54, 21 (1996).
5. T. Kinoshita, Phys. Rev. Lett. 47, 1573 (1981); Review of Particle Properties, Phys. Rev. D 54, 21 (1996).
6. V. Barger and R. Philips, "Collider Physics", Addison-Wesley (Menlo-Park, 1987).
7. Review of Particle Properties, Phys. Rev. D 54, 222 (1996).
8. Junichi Kanzaki, in Proceedings of the second workshop on JLC, ed. by S. Kawabata

(KEK 91-10).

9. H. Georgi, Phys. Lett. B 298, 187 (1993); R. S. Chivukula and E. H. Simmons, Phys. Lett. B 388, 788 (1996).

Figure Captions

Fig. 1. The cross-section for the process $e^+e^- \rightarrow HZ$ for $m_H = 150$ Gev plotted against $\sqrt{s}$.

(a) constructive interference (b) standard model and (c) destructive interference.

Fig. 2 Same as Fig. 1 with $m_H = 200$ Gev.